

## Effect of cation symmetry on the low-frequency spectra of imidazolium ionic liquids: OKE and Raman spectroscopic measurements and DFT calculations


Dong Xiao<sup>a</sup>, Larry G. Hines Jr.<sup>a</sup>, Mark W. Holtz<sup>b</sup>, Kihyung Song<sup>c</sup>, Richard A. Bartsch<sup>a</sup>, Edward L. Quitevis<sup>a,b,\*</sup>

<sup>a</sup> Department of Chemistry and Biochemistry, Texas Tech University, Lubbock, TX 79409, United States

<sup>b</sup> Department of Physics, Texas Tech University, Lubbock, TX 79409, United States


<sup>c</sup> Department of Chemistry, Korea National University of Education, Chongwon, Chungbuk 363-391, Republic of Korea

# Principle of optical kerr effect



# OHD-RIKE and DFT(density functional theory)




- OHD-RIKE



- DFT


Quantum mechanical modeling method used in physics and chemistry to investigate the electronic structure (principally the ground state) of many-body systems, in particular atoms, molecules, and the condensed phases.

# Sample




# Raman Spectra



- Polarized Raman spectra





- Depolarized Raman spectra



## Reduced Spectral Density by Kerr signal



# Calculated Raman spectra



region A1: 0 < w ≤ 100 /cm

region A2: 100 < w ≤ 125 /cm

region A3: 125 < w ≤ 200 /cm

## Concluding remarks

- According to Kerr experiment on two kinds of ionic liquids, the intermolecular motion could be determined by symmetry of cation.
- In the range from 100 to 200 /cm, a weak peak originate from the anion.
- The intramolecular motion should be considered besides the intermolecular motion when you do peak assignment.